

B1

PCT/EP 01 / 06 4 33




ÖSTERREICHISCHES PATENTAMT

A-1014 WIEN, KOHLMARKT 8 - 10

REC'D 13 SEP 2001

WIPO

PCT



Aktenzeichen A 1973/2000

Das Österreichische Patentamt bestätigt, dass

die Firma CISTEM BIOTECHNOLOGIES GMBH
in A-1030 Wien, Rennweg 95B,

am **23. November 2000** eine Patentanmeldung betreffend

"Immunstimulatorische Oligodesoxynukleotide",

überreicht hat und dass die beigeheftete Beschreibung samt Zeichnungen
mit der ursprünglichen, zugleich mit dieser Patentanmeldung überreichten
Beschreibung samt Zeichnungen übereinstimmt.

Österreichisches Patentamt

Wien, am 12. April 2001

Der Präsident:

i. A.




**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

11111111



ÖSTERREICHISCHES PATENTAMT

Verwaltungsstellen-Direktion

....200.- s. 58,14... e

Kanzleigebühr bezahlt.

Baraoun

A1973/2000

R 37332

(51) Int. Cl. :

Urtext

AT PATENTSCHRIFT

(11) Nr.

(73) Patentinhaber: CISTEM BIOTECHNOLOGIES GMBH
Wien (AT)

(54) Gegenstand: Immunstimulatorische Oligodesoxynukleotide

(61) Zusatz zu Patent Nr.

(67) Umwandlung aus GM

(62) Ausscheidung aus :

(22) (21) Angemeldet am: 23. NOV. 2000

(33) (32) (31) Unionspriorität :

(42) Beginn der Patentdauer:

Längste mögliche Dauer:

(45) Ausgegeben am :

(72) Erfinder :

(60) Abhängigkeit:

(56) Entgegenhaltungen, die für die Beurteilung der Patentierbarkeit in Betracht gezogen wurden:

03370

- 1 -

The present invention relates to immunostimulatory oligodeoxynucleic molecules (ODNs) and pharmaceutical compositions containing such ODNs.

Vaccines can save more lives (and resources) than any other medical intervention (Nossal, 1998). Owing to world-wide vaccination programs the incidence of many fatal diseases has been decreased drastically. Although this notion is valid for a whole panel of diseases, e.g. tuberculosis, diphtheria, pertussis, measles and tetanus, there are no effective vaccines for numerous infectious disease including most viral infections, such as AIDS. There are also no effective vaccines for other diseases, infectious or non-infectious claiming millions the lives of millions of patients per year including malaria or cancer. In addition, the rapid emergence of antibiotic-resistant bacteria and microorganisms calls for alternative treatments with vaccines being a logical choice. Finally, the great need for vaccines is also illustrated by the fact that infectious diseases, rather than cardiovascular disorders or cancer or injuries remain the largest cause of death and disability in the world (Bloom and Widdus, 1998).

From an immunological point of view one major problem in the field of vaccines today is that traditional vaccines (and/or the immune-modulating compounds contained within these preparations) are designed to induce high levels of antibodies (Harrow and Lane, 1988). However, antibodies on their own are not effective in preventing a large number of diseases including most illnesses caused by viruses, intracellular bacteria, certain parasites and cancer. Examples for such diseases are, but are not restricted to, the above-mentioned HIV virus or Plasmodium spec. in case of malaria. In numerous experimental systems it has been shown that the cellular arm of the immune system, including T cells, rather than the humoral arm, is important for these indications. Therefore, novel, innovative technologies are needed to overcome the limitations of conventional vaccines. The focus must be on technologies that reliably induce the cellular immune system, including antigen specific T cells, which recognize molecules expressed

03377

- 2 -

from normal cells and, simultaneously, antibodies secreted by B cells recognising pathogens in extracellular compartments.

Several established vaccines consist of live attenuated organism where the risk of reversion to the virulent wild-type strain exists. In particular in immunocompromised hosts this can be a live threatening scenario. Alternatively, vaccines are administered as a combination of pathogen-derived antigens together with compounds that induce or enhance immune responses against these antigens (these compounds are commonly termed adjuvant), since these subunit vaccines on their own are generally not effective.

Whilst there is no doubt that the above vaccines are valuable medical treatments, there is the disadvantage that, due to their complexity, severe side effects can be evoked, e.g. to antigens that are contained in the vaccine that display cross-reactivity with molecules expressed by cells of vaccinated individuals. In addition, existing requirements from regulatory authorities, e.g. the World Health Organization (WHO), the Food and Drug Administration (FDA), and their European counterparts, for exact specification of vaccine composition and mechanisms of induction of immunity, are difficult to meet.

Antigen presenting cells belong to the innate immune system, which has evolved as a first line host defence that limits infection early after exposure to microorganisms (Hoffmann et al., 1999). Cells of the innate immune system recognize patterns or relatively non-specific structures expressed on their targets rather than more sophisticated, specific structures which are recognised by the adaptive immune system (Hoffmann et al., 1999). Examples of cells of the innate immune system are macrophages and dendritic cells but also granulocytes (e.g. neutrophils), natural killer cells and others. By contrast, cells of the adaptive immune system recognize specific, antigenic structures, including peptides, in the case of T cells and peptides as well as three-dimensional structures in the case of B cells. The adaptive immune system is much more specific and sophisticated than the innate immune system and improves upon repeat exposure to a given pathogen/antigen. Phylogenetically, the innate immune system is

03370

- 3 -

Nevertheless, the innate immune system is critical during the initial phase of antigenic exposure since, in addition to containing pathogens, cells of the innate immune system, i.e. APCs, prime cells of the adaptive immune system and thus trigger specific immune responses leading to clearance of the intruders. In sum, cells of the innate immune system and in particular APCs play a critical role during the induction phase of immune responses by a) containing infections by means of a primitive pattern recognition system and b) priming cells of the adaptive immune system leading to specific immune responses and memory resulting in clearance of intruding pathogens or of other targets (Roitt et al., 1998). These mechanisms may also be important to clear or contain tumor cells.

As mentioned above, cells of the innate immune system recognise patterns expressed on their respective targets. Examples are lipopolysaccharides (LPS) in the case of Gram-negative bacteria, mycobacterial glycolipids, lipoteichoic acids of Gram-positive bacteria, mannans of yeast and double stranded RNAs of viruses (Hoffmann et al., 1999). In addition they may recognise patterns such as altered glycosylations of proteins on tumor cells.

Recent findings describe DNAs of protozoan or lower eukaryotes as a further pattern recognised by the innate (but possibly also by the adaptive) immune system of mammals (and probably most if not all vertebrates) (Krieg, 1996; Lipford et al., 1998).

The immune system recognises lower organisms including bacteria probably due to structural and sequence usage differences between pathogen and host DNA. In particular short stretches of DNA, derived from non-vertebrates or in form of short synthetic ODNs containing nonmethylated cytosine-guanine dinucleotides (CpG) in a certain base context, are targeted (Krieg et al., 1995). CpG motifs are found at the expected frequency in bacterial DNA but are much less frequent in vertebrate DNA (Lipford et al., 1998; Pisetky, 1999). In addition, non-vertebrate (i.e. bacterial) CpG motifs are not methylated whereas vertebrate CpG sequences are. These differences between bacterial DNA and verte-

03370

- 4 -

Natural CpG-containing DNA, ODNs, as well as thiophosphate-substituted (exchange of thiophosphate residues for phosphate) ODNs containing CpG motifs (CpG-ODN) are not only potent activators of immune cell proliferation and humoral immune responses (Krieg et al., 1995), but also stimulate strong cellular immune responses (reviewed in Lipford et al., 1998). DNA/ODNs containing non-methylated CpG motifs can directly activate monocytic cells (dendritic cells, macrophages) and B cells. Likely, natural killer (NK) cells are not directly activated but respond to monocyte-derived IL-12 (interleukin 12) with a marked increase in their IFN- γ production (Chace et al., 1997). In consequence, the induction of monocytes and NK cells by CpG DNA promotes the induction of Th1-type responses and the development of cytotoxic T cells.

Ribonucleic acid based on inosine and cytosine, like polyinosinic-polycytidylic acid (poly I:C), is known to promote Th1-specific immune responses. It is known to stimulate macrophages to produce cytokines such as IL-1 α and IL-12 (Manetti et al., 1995), it is also known as a potent interferon type 1 inducer (Manetti et al., 1995) and a potent NK cell stimulator (Cavanaugh et al., 1996).

This effect, however, was strictly restricted to ribonucleic acid containing inosine and cytidine residues (WO98/16247).

Investigations by the inventors of the present invention showed that ODNs containing non-methylated CpG motifs, although being efficient in stimulating immune system, have essential disadvantages, especially with respect to specificity (high background) and induction of side effects, such as high systemic TNF- α generation. High systemic TNF- α release is known to cause toxic shock syndrome, which can cause death of afflicted patients.


It is therefore an object of the present invention to provide suitable novel ODNs which do not have such drastic side effects as ODNs based on CpG sequences. It is a further object to reduce the side effects of pharmaceutical compositions containing known

033070

- 5 -

tical compositions with efficient, immunostimulatory properties which are suitable for vaccination of animals, especially of mammals, including humans.

This object is solved by immunostimulatory oligodeoxynucleic acid molecule (ODN) having the structure according to formula (I)



(I),

any X is O or S,

wherein

any NMP is a 2' deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine-, deoxyguanosine-, deoxyinosine-, deoxycytosine-, deoxyuridine-, deoxythymidine-, 2-methyl-deoxyinosine-, 5-methyl-deoxycytosine-, deoxypseudouridine-, deoxyribosepurine-, 2-amino-deoxyribosepurine-, 6-S-deoxyguanine-, 2-dimethyl-deoxyguanosine- or N-isopentenyl-deoxyadenosine-monophosphate or -monothiophosphat,

NUC is a 2' deoxynucleoside, selected from the group consisting of deoxyadenosine-, deoxyguanosine-, deoxyinosine-, deoxycytosine-, deoxyuridine-, deoxythymidine-, 2-methyl-deoxyinosine-, 5-methyl-deoxycytosine-, deoxypseudouridine-, deoxyribosepurine-, 2-amino-deoxyribosepurine-, 6-S-deoxyguanine-, 2-dimethyl-deoxyguanosine- or N-isopentenyl-deoxyadenosine,

a and b are integers from 0 to 100 with the proviso that a + b is between 4 and 150,

B and E are common groups for 5' or 3' ends of nucleic acid molecules.

000000

- 6 -

Surprisingly it turned out that ODNs containing deoxyinosine residues (I-ODNs) show an immunostimulatory effect comparable or in many instances even better than ODNs containing CpG motifs. Moreover, ODNs according to the present invention produce more specific immune responses to a given antigen or antigen fragment than CpG ODNs. In addition, ODNs according to the present invention reduced the induction of adverse side reactions, especially the induction of systemic TNF- α or IL-6.

Whereas certain immunostimulatory effects had been described for inosine containing RNA molecules, such as poly-IC or the molecules mentioned in WO98/16247, it surprisingly turned out that deoxynucleic acid molecules containing deoxyinosine residues, may be good immunostimulating ODNs.

In addition, the I-ODNs according to the present invention are - in contrast to ODNs based on the specific CpG motif - not dependent on a specific motif or a palindromic sequence as described for the CpG oligonucleotides (see e.g. EP 0 468 520 A2, WO96/02555, WO98/18810, WO98/37919, WO98/40100, WO98/52581, WO99/51259 and WO99/56755, all incorporated herein by reference). Therefore, one group of I-ODNs according to the present invention may preferably contain a CI motif (and therefore ODNs described in these incorporated references, wherein one or more guanosine residues are replaced with deoxyinosine residues are preferred embodiments of the present ODNs). It is not necessary for its principle immunostimulatory property, since I-ODNs with an Inosine not placed in a CI or IC context exhibit immunostimulatory properties as well.

The I-ODN according to the present invention is therefore a DNA molecule containing a deoxyinosine residue which is preferably provided in single stranded form.

The I-ODN according to the present invention may be isolated through recombinant methods or chemically synthesized. In the latter case, the I-ODN according to the present invention may also contain modified oligonucleotides which may be synthesized using standard chemical transformations, such as methylphosphon-

03370

- 7 -

as phosphotriesters, phosphoamidates and phosphorodithiorates. Other non-phosphorous based modified oligonucleotides can also be used (Stirchak et al., MAR 17 (1989), 6129-6141), however, mono-phosphates or monothiophosphates being the preferred 2'deoxy nucleoside monophosphate to be used in the present invention.

The NMPs of the I-ODNs according to the present invention are preferably selected from the group consisting of deoxyadenosine-, deoxyguanosine-, deoxyinosine-, deoxycytosine-, deoxyuridine-, deoxythymidine-, 2-methyl-deoxyinosine-, 5-methyl-deoxycytosine-monophosphate or -monothiophosphate (as usual, the phosphate or thiophosphate group is 5' of the deoxyribose). Whereas it is essential for the ODNs based on the CpG motif that this motif is unmethylated, this is surprisingly not the case for the ODNs according to the present invention, wherein e.g. 2-methyl-deoxyinosine or 5-methyl-deoxycytosine residues have no general negative effect on immunostimulatory properties of the ODNs according to the present invention. Alternatively, instead of the 2-deoxy-forms of the NMPs, also other, especially inert, groups may be present at the 2-site of the ribose group, such as e.g. -F, -NH₂, -CH₃, especially -CH₃. Of course, -OH and SH groups are excluded for the I-ODNs according to the present invention to be present on the 2'-site of the ribose, especially the ribose residue for the inosine NMP.

The length of the ODNs according to the present invention is in the range of the standard ODNs used according to the prior art. Therefore molecules with a total length under 4 and above 150 show gradually decreasing immunostimulatory potential. Preferred ODNs contain between 10 and 60, especially between 15 and 40 bases (nucleosides), implying that a + b in formula I is between 10 and 60, preferably between 15 and 40 in these preferred embodiments.

Whereas the ribonucleic acid molecules containing inosine and cytidine described to be immunostimulatory in the prior art have been large and relatively undefined polynucleic acids with molecular weights far above 200,000 (a commercially available poly-inosinic-polycytidylic acid from Sigma Chemicals has a molecular

03370

- 8 -

residues). The molecules according to the present invention are DNA molecules of much shorter length with a well defined length and composition, being highly reproducible in products.

It is further preferred that the deoxyinosine containing NMP of the I-ODNs according to formula I is a monothiophosphate with one to four sulfur atoms and that also further NMPs, especially all further NMPs, are present as nucleoside monothiophosphates, because such ODNs display higher nuclease resistance (it is clear for the present invention that the "mono" in the "monothiophosphates" relates to the phosphate, i.e. that one phosphate group (one phosphor atom) is present in each NMP). Preferably, at least one of X_1 and X_2 is S and at least one of X_3 and X_4 is O in the NMPs according to the present invention. Preferably, X_3 and X_4 are O. (X_3 may be (due to synthesis of the NMP) derived e.g. from the phosphate group or from the 3'-group of the NMP-ribose).

Preferably the ODNs according to the present invention contain the sequence

nhh hhh wdi nhh hhh hhh wn,
 nhh wdi din hhh hdi ndi nh,
 nhh hhh wdi dhh hhh hhh wn or
 nhh wdi did hhh hdi ddi dh,

wherein

any n is a 2'-deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine-, deoxyguanosine-, deoxycytosine- or deoxythymidine-monophosphate or -monothiophosphate,

any h is a 2'-deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine-, deoxycytosine- or deoxythymidine-monophosphate or -monothiophosphate

i is deoxyinosine-monophosphate or -monothiophosphate,

any w is a 2'-deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine- or deoxythymidine-monophosphate or -monothiophosphate, and

any d is a 2'-deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine-, deoxyguanosine- or deoxythymidine-monophosphate or -monothiophosphate.

033270

- 9 -

is not necessary for the I-ODNs according to the present invention. However, ODNs containing a CI motif are preferred so that in a preferred embodiment the ODN according to formula I contains at least one 2'-deoxycytosine-monophosphate or -monothiophosphate 3'-adjacent to a 2'-deoxyinosine-monophosphate or -monothiophosphate to form such a 5'-CI 3'-motif.

Preferred ODNs according to the present invention contain one or more of the sequence

gacitt,
iacitt,
gaictt,
iaictt,

wherein

a is deoxyadenosine-monophosphate or -monothiophosphate,
g is deoxyguanosine-monophosphate or -monothiophosphate,
i is deoxyinosine-monophosphate or -monothiophosphate,
c is deoxycytosine-monophosphate or -monothiophosphate and
t is deoxythymidine-monophosphate or -monothiophosphate.

The I-ODNs according to the present invention are especially suitable for application in the pharmaceutical field, e.g. to be applied as a medicine to an animal or to humans. They are specifically adapted to act as an immunostimulatory agent, especially in or together with vaccine compositions.

Therefore, the present invention also relates to a pharmaceutical composition comprising an ODN according to the present invention.

Since a preferred pharmaceutical composition according to the present invention is a vaccine, this composition should contain an antigen besides the ODN according to the present invention. The potential of this antigen to raise a protection/immune response of the vaccinated individual is strongly increased by combining it with the ODNs according to the present invention, especially due to their immunostimulatory activity.

A vaccine can contain a whole variety of different antigens. Examples of antigens are whole killed organisms such as inactivated

03070

- 10 -

gens may also consist of subfractions of these organisms/tissues, of proteins, or, in their most simple form, of peptides. Antigens can also be recognised by the immune system in form of glycosylated proteins or peptides and may also be or contain polysaccharides or lipids. Short peptides can be used since for example cytotoxic T cells (CTL) recognize antigens in form of short usually 8-11 amino acids long peptides in conjunction with major histocompatibility complex (MHC) (Rammensee et al., Immunogenetics 41, (1995), 178-228). B cells recognize longer peptides starting at around 15 amino acids (Harrow et al, Cold Spring Harbor: Cold Spring Harbor Laboratory, (1988)). By contrast to T cell epitopes, the three dimensional structure of B cell antigens may also be important for recognition by antibodies. In order to obtain sustained, antigen-specific immune responses, adjuvants are helpful to trigger immune cascades that involve all cells of the immune system necessary. Primarily, adjuvants are acting, but are not restricted in their mode of action, on so-called antigen presenting cells (APCs). These cells usually first encounter the antigen(s) followed by presentation of processed or unmodified antigen to immune effector. Intermediate cell types may also be involved. Only effector cells with the appropriate specificity are activated in a productive immune response. The adjuvant may also locally retain antigens and co-injected other factors. In addition the adjuvant may act as a chemoattractant for other immune cells or may act locally and/or systemically as a stimulating agent for the immune system.

The antigens to be used in the present compositions are not critical. Preferably, proteins or peptides derived from a viral or a bacterial pathogen or from fungi or parasites are used as such antigens (including derivatized antigens or glycosylated or lipidated antigens or polysaccharides or lipids). Another preferred source of antigens are tumor antigens. Preferred pathogens are selected from human immunodeficiency virus (HIV), hepatitis A and B viruses, hepatitis C virus (HCV), rous sarcoma virus (RSV), Epstein Barr virus (EBV) Influenza virus, Rotavirus, Staphylococcus aureus, Chlamydia pneumonias, Chlamydia trachomatis, Mycobacterium tuberculosis, Streptococcus pneumonias, Bacillus anthracis, Vibrio cholerae. Plasmodium sp. (Pl. falciparum, Pl.

03370

- 11 -

also be molecules expressed by cancer cells (tumor antigens). The derivation process may include the purification of a specific protein from the pathogen/cancer cells, the inactivation of the pathogen as well as the proteolytic or chemical derivatization or stabilisation of such a protein. In the same way also tumor antigens (cancer vaccines) or autoimmune antigens may be used in the pharmaceutical composition according to the present invention. With such compositions a tumor vaccination or a treatment for autoimmune diseases may be performed.

In the case of peptide antigens the use of peptide mimitopes/agonists/superagonists/antagonists or peptides changed in certain positions without affecting the immunologic properties or non-peptide mimitopes/agonists/superagonists/antagonists (reviewed in Sparbier and Walden, 1999) is included in the current invention. Peptide antigens may also contain elongations either at the carboxy or at the amino terminus of the peptide antigen facilitating interaction with the polycationic compound(s) or the immunostimulatory compound(s). For the treatment of autoimmune diseases peptide antagonists may be applied.

Antigens may also be derivatized to include molecules enhancing antigen presentation and targeting of antigens to antigen presenting cells.

In one embodiment of the invention the pharmaceutical composition serves to confer tolerance to proteins or protein fragments and peptides which are involved in autoimmune diseases. Antigens used in this embodiments serve to tolerize the immune system or down-regulate immune responses against epitopes involved in autoimmune processes.

Preferably the pharmaceutical composition according to the present invention, especially in the form of a vaccine, further comprises a polycationic polymer, preferably a polycationic peptide, especially polyarginine, polylysine or an antimicrobial peptide.

The polycationic compound(s) to be used according to the present

00370

- 12 -

tionic compounds are selected from basic polypeptides, organic polycations, basic polyaminoacids or mixtures thereof. These polyaminoacids should have a chain length of at least 4 amino acid residues (see: Tuftsin as described in Goldman et al (1983)). Especially preferred are substances containing peptidic bounds, like polylysine, polyarginine and polypeptides containing more than 20%, especially more than 50% of basic amino acids in a range of more than 8, especially more than 20, amino acid residues or mixtures thereof. Other preferred polycations and their pharmaceutical compositions are described in WO 97/30721 (e.g. polyethyleneimine) and WO 99/38528. Preferably these polypeptides contain between 20 and 500 amino acid residues, especially between 30 and 200 residues.

These polycationic compounds may be produced chemically or recombinantly or may be derived from natural sources.

Cationic (poly)peptides may also be polycationic anti-bacterial microbial peptides with properties as reviewed in (Ganz and Lehrer, 1999; Hancock, 1999). These (poly)peptides may be of prokaryotic or animal or plant origin or may be produced chemically or recombinantly (Andreu and Rivas, 1998; Ganz and Lehrer, 1999; Simmaco et al., 1998). Peptides may also belong to the class of defensins (Ganz, 1999; Ganz and Lehrer, 1999). Sequences of such peptides can be, for example, be found in the Antimicrobial Sequences Database under the following internet address:

<http://www.bbcm.univ.trieste.it/~tossi/pag1.html>

Such host defense peptides or defensives are also a preferred form of the polycationic polymer according to the present invention. Generally, a compound allowing as an end product activation (or down-regulation) of the adaptive immune system, preferably mediated by APCs (including dendritic cells) is used as polycationic polymer.

Especially preferred for use as polycationic substance in the present invention are cathelicidin derived antimicrobial peptides or derivatives thereof (A 1416/2000, incorporated herein by reference), especially antimicrobial peptides derived from mammal

03370

- 13 -

tive compounds, such as (human) growth hormone.

Polycationic compounds derived from natural sources include HIV-REV or HIV-TAT (derived cationic peptides, antennapedia peptides, chitosan or other derivatives of chitin) or other peptides derived from these peptides or proteins by biochemical or recombinant production. Other preferred polycationic compounds are cathelin or related or derived substances from cathelin. For example, mouse cathelin is a peptide which has the amino acid sequence $\text{NH}_2\text{-RLAGLLRKGGEKIGEKLLKIGOKIKNFFQKLVPQPE-COOH}$. Related or derived cathelin substances contain the whole or parts of the cathelin sequence with at least 15-20 amino acid residues. Derivations may include the substitution or modification of the natural amino acids by amino acids which are not among the 20 standard amino acids. Moreover, further cationic residues may be introduced into such cathelin molecules. These cathelin molecules are preferred to be combined with the antigen and the immunogenic ODN according to the present invention. However, these cathelin molecules surprisingly have turned out to be also effective as an adjuvant for a antigen without the addition of further adjuvants. It is therefore possible to use such cathelin molecules as efficient adjuvants in vaccine formulations with or without further immunactivating substances.

Another preferred polycationic substance to be used according to the present invention is a synthetic peptide containing at least 2 KLK-motifs separated by a linker of 3 to 7 hydrophobic amino acids (A 1789/2000, incorporated herein by reference).

It was very surprising that the immunostimulating effect of the pharmaceutical composition according to the present invention was significantly higher than it could be expected from the addition of the effects of each single component or even the addition of the effects of the ODN or the polycation with the antigen.

B and E in formula I are common groups for 5' and/or 3' ends of nucleic acid molecules. Examples for such groups are readily available for the skilled man in the art (see e.g. "Oligonucleo-

03370

- 14 -

invention B and/or E are preferably selected independently from -H, -CH₃, -COCH₃, -OH, -CHO, a phosphate, thiophosphate, sulfate or a thiosulfate group, or a phosphoalkylgroup, especially with an alkyl length of C₁-C₆ and/or with a terminal amino group (the amino group may e.g. be used for further labelling of the I-ODNs according to the present invention, e.g. -PO₄-(CH₂)_n-NH₂ or -PO₄-(CH₂)_n-NH-Label). Especially preferred as B are nucleosides, especially the 2'-deoxynucleotides mentioned above (i.e. without the phosphate or thiophosphate group). Alternatively these groups may also contain linker groups to other molecules, especially carrier molecules or labels. In such forms of ODNs wherein the ODNs are bound to solid surfaces or particles or labels, these surfaces, particles, labels, etc. are then also part of the B and/or E groups.

Of course, any ionised (salt) form or tautomeric forms of the molecules according to formula I are included in this formula I.

The pharmaceutical composition according to the present invention may further comprise further active ingredients (pharmaceutically active substances), especially substances which are usable in a vaccine connection. Preferred embodiments of such further active ingredients are cytokines, antiinflammatory substances, antimicrobial substances or combinations thereof.

Of course, the pharmaceutical composition according to the present invention may further contain auxiliary substances, especially a pharmaceutically acceptable carrier, buffer substances, stabilizers or combinations thereof.

The relative amounts of the ingredients in the present pharmaceutical composition are highly dependent on the necessities of the individual antigen and on the animal/human to which this composition should be applied to. Therefore, the pharmaceutical composition according to the present invention preferably contains one or more ODNs according to the present invention, preferably 1 pg to 10 g, preferably 1 ng to 1 g, more preferred 100 ng to 10 mg, especially 10 µg to 1 mg. The antigen as well as the polycationic polymer may be applied in similar dosages, a range of 1 to 10.000

03370

- 15 -

ferred.

The present compositions may be applied to a patient, e.g. a vaccination candidate, in efficient amounts e.g. by weekly, bi-weekly or monthly intervals. Patients to be treated with the present compositions may also be vaccinated repeatedly or only once. A preferred use of the present invention is the active immunisation, especially of humans or animals without protection against the specific antigen.

The route of application for the present composition is not critical, e.g. subcutaneous, intramuscular, intradermal or transdermal injection is suitable as well as oral uptake.

It is also possible to apply the present composition separately e.g. by injecting the immunostimulating substance separately from the antigen/polycation composition. The present invention is therefore also directed to a kit comprising a composition containing the antigen and the polycationic polymer as one component and a composition containing the immunostimulating or chemotactic substance as a second component.

The components may be applied at the same site or time, however, an application at different sites or at a different time or for a different time period is also possible. It is also possible to vary the systemic or local applications of the composition or the components, respectively.

Details of the present invention are described by the following examples and the figures, but the invention is of course not limited thereto.

Fig. 1 shows the immune response against the ovalbumin-derived peptide OVA₂₅₇₋₂₆₄ after the injection of OVA₂₅₇₋₂₆₄, poly-L-arginine (pR 60) and deoxyinosine I-containing oligodeoxynucleotides I-ODN) or CpG 1668. Mice were injected into the hind footpads with mixtures as indicated. Four days later draining lymph node cells were ex vivo stimulated with OVA₂₅₇₋₂₆₄, an irrelevant peptide.

03370

- 16 -

ing cells was determined 24 hours later using an ELISPOT assay. Results are expressed as the number of spots/ 1×10^6 lymph node cells with standard deviation of triplicates.

Fig. 2 shows the induction of systemic TNF- α production after the injection of OVA₂₅₇₋₂₆₄, poly-L-arginine (pR 60) and I-containing oligodeoxynucleotides (I-ODN) or CpG 1668. Mice were injected into the hind footpads with mixtures as indicated. One hour after injection blood was taken from the tail vein and serum was prepared. The concentration of TNF- α in the sera was determined using an ELISA.

Fig.3 shows the immune response against the Ovalbumin-derived peptide OVA₂₅₇₋₂₆₄ after the injection of OVA₂₅₇₋₂₆₄, poly-L-arginine (pR60) and deoxyinosine -containing oligodeoxynucleotides (I-ODN) , CpG 1668 or GpC. Mice were injected into the hind footpads with mixtures as indicated. Four days later, draining lymph node cells were ex vivo stimulated with OVA₂₅₇₋₂₆₄, an irrelevant peptide mTRP2₁₈₁₋₁₈₈ (murine tyrosinase related protein-2, VYDFFVWL), pR 60. The number of IFN- γ producing cells was determined 24 hours later using an ELISPOT assay. Results are expressed as the number of spots/ 1×10^6 lymph node cells with standard deviation of triplicates.

Fig. 4 shows the induction of systemic TNF- α production after the injection of OVA₂₅₇₋₂₆₄, poly-L-arginine (pR 60) and I-containing oligodeoxynucleotides (I-ODN), GpC or CpG 1668. Mice were injected into the hind footpads with mixtures as indicated. One hour after injection blood was taken from the tail vein and serum was prepared. The concentration of TNF- α and IL-6 in the sera was determined using cytokin-specific ELISAs.

Fig. 5 shows the immune response against the Ovalbumin-derived peptide OVA₂₅₇₋₂₆₄ after the injection of TRP-2, poly-L-arginine, CpG 1668 or random sequences containing deoxyinosine. Mice were injected into the hind footpads with mixtures as indicated. Four days later, draining lymph node cells were ex vivo stimulated

03370

- 17 -

IFN- γ producing cells was determined 24 hours later using an ELISPOT assay. Results are expressed as the number of spots/ 1×10^6 lymph node cells with standard deviation of triplicates.

EXAMPLES

In all experiments thiophosphate-substituted ODNs (with thiophosphate residues substituting for phosphate, hereafter called "thiophosphate substituted oligodeoxynucleotides") were used since such ODNs display higher nuclease resistance (Ballas et al., 1996; Krieg et al., 1995; Parronchi et al., 1999).

Example 1

The combined injection of I-ODN and poly-L-arginine (pR 60) synergistically enhances the immune response against an Ovalbumin-derived peptide.

| | |
|--------------------------|---|
| Mice | C57BI/6 (Harlan/Olac) |
| Peptide | OVA ₂₅₇₋₂₆₄ -Peptide (SIINFEKL), a MHC class I (H-2Kb)-restricted epitope of chicken ovalbumin (Rotzschke et al., 1991), was synthesized using standard solid phase F-moc chemistry synthesis, HPLC purified and analysed by mass spectroscopy for purity. |
| | Dose: 300 μ g/mouse |
| Poly-L-arginine60 (pR60) | Poly-L-arginine with an average degree of polymerization of 60 arginine residues; SIGMA chemicals |
| | Dose: 100 μ g/mouse |
| CpG-ODN 1668 | thiophosphate substituted ODNs containing a CpG motif: tcc atg <u>acg</u> ttc ctg atg ct, were synthesized by NAPS GmbH, Göttingen. |
| | Dose: 5 nmol/mouse |

03377

- 18 -

I-ODN 1

thiophosphate substituted ODNs
containing Inosine:
tcc ati aci ttc ctg atg ct, were
synthesized by NAPS GmbH, Göttin-
gen.

Dose: 5nmol/mouse

I-ODN 2

thiophosphate substituted ODNs
containing Inosine:
tcc atg aci ttc ctg atg ct, were
synthesized by NAPS GmbH, Göttin-
gen.

Dose: 5nmol/mouse

I-ODN 3

thiophosphate substituted ODNs
containing Inosine:
tcc ati aci ttc cti ati ct, were
synthesized by NAPS GmbH, Göttin-
gen.

Dose: 5nmol/mouse

Experimental groups (5 mice per group)

1. OVA₂₅₇₋₂₆₄
2. OVA₂₅₇₋₂₆₄ + pR 60
3. OVA₂₅₇₋₂₆₄ + CpG 1668
4. OVA₂₅₇₋₂₆₄ + I-ODN 1
5. OVA₂₅₇₋₂₆₄ + I-ODN 2
6. OVA₂₅₇₋₂₆₄ + I-ODN 3
7. OVA₂₅₇₋₂₆₄ + CpG 1668 + pR 60
8. OVA₂₅₇₋₂₆₄ + I-ODN 1 + pR 60
9. OVA₂₅₇₋₂₆₄ + I-ODN 2 + pR 60
10. OVA₂₅₇₋₂₆₄ + I-ODN 3 + pR 60

On day 0 mice were injected into each hind footpad with a total volume of 100 µl (50 µl per footpad) containing the above mentioned compounds. Animals were sacrificed 4 days after injection and popliteal lymph nodes were harvested. Lymph nodes were passed through a 70 µm cell strainer and washed twice with DMEM medium (GIBCO BRL) containing 5% fetal calf serum (FCS, SIGMA

03370

- 19 -

scribed (Miyahira et al., 1995). This method is a widely used procedure allowing the quantification of antigen-specific T cells. Lymphocytes were stimulated ex vivo in triplicates with medium background-control, OVA₂₅₇₋₂₆₄-peptide alone, or by combinations of OVA-peptide with pR 60, I-ODN1 or I-ODN2 or I-ODN3 and Concanavalin A (Con A). Spots representing single IFN- γ producing T cells were counted and the number of background spots was subtracted from all samples. There were many spots detected after the stimulation with Con A (data not shown) indicating a good condition of the used lymphocytes. For each experimental group of mice the number of spots/ 1×10^6 cells are illustrated in Figure 1, the standard deviation of ex vivo-stimulated triplicates are given.

One hour after injection blood was taken from the tail vein and serum was prepared to determine the induction of systemic TNF- α using an ELISA (Figure 2).

Example 2

The exchange of Guanosine by desoxy-Inosine converts the non-immunogeneic GpC-sequence to a highly immunogeneic one, especially when combined with poly-L-arginine (pR60).

| | |
|---------------------------|--|
| Mice | C57Bl/6 (Harlan/Olac) |
| Peptide | OVA ₂₅₇₋₂₆₄ -Peptide (SIINFEKL), a MHC class I (2Kb)-restricted epitope of chicken ovalbumin (Rotzschke et al., 1991), was synthesized using standard solid phase F-moc synthesis, HPLC purified and analysed by mass spectroscopy for purity. Dose: 300 μ g/mouse |
| Poly-L-arginine 60 (pR60) | Poly-L-arginine with an average degree of polymerization of 60 arginine residues; SIGMA chemicals Dose: 100 μ g/mouse |

03370

- 20 -

- CpG-ODN 1668 thiophosphate substituted ODNs containing a CpG motif: tcc atg acg ttc ctg atg ct, were synthesized by NAPS GmbH, Göttingen.
Dose: 5nmol/mouse
- GpC-ODN thiophosphate substituted ODNs containing an non-immunogeneic GpC motif: tcc atg agc ttc ctg atg ct were synthesized by NAPS GmbH, Göttingen.
Dose: 5nmol/mouse
- I-ODN 9 thiophosphate substituted ODNs containing Inosine: tcc atg aic ttc ctg atg ct were synthesized by NAPS GmbH, Göttingen.
Dose: 5nmol/mouse
- I-ODN 10 thiophosphate substituted ODNs containing Inosine: tcc ati aic ttc cti ati ct were synthesized by NAPS GmbH, Göttingen.
Dose: 5nmol/mouse

Experimental groups (5 mice per group)

1. OVA₂₅₇₋₂₆₄
2. OVA₂₅₇₋₂₆₄ + pR 60
3. OVA₂₅₇₋₂₆₄ + CpG 1668
4. OVA₂₅₇₋₂₆₄ + GpC
5. OVA₂₅₇₋₂₆₄ + I-ODN 9
6. OVA₂₅₇₋₂₆₄ + I-ODN 10
7. OVA₂₅₇₋₂₆₄ + CpG 1668 + pR 60
8. OVA₂₅₇₋₂₆₄ + GpC + pR 60
9. OVA₂₅₇₋₂₆₄ + I-ODN 9 + pR 60
10. OVA₂₅₇₋₂₆₄ + I-ODN 10 + pR 60

On day 0 mice were injected into each hind footpad with a total volume of 100µl (50µl per footpad) containing the above mentioned compounds. Animals were sacrificed 4 days after injection and popliteal lymph nodes were harvested. Lymph nodes were passed through a 70µm cell strainer and washed twice with DMEM medium (GIBCO BRL) containing 5% fetal calf serum (FCS, SIGMA chemicals). Cells were adjusted to 3x10⁶ cells/ml in DMEM/5%FCS. An IFN-γ ELISPOT assay was carried out in triplicates as described (Miyahira et al., 1995). This method is a widely used procedure allowing the quantification of antigen-specific T

00070

- 21 -

medium (background), OVA₂₅₇₋₂₆₄-peptide, an irrelevant peptide mTRP-2₁₈₁₋₁₈₈ (murine tyrosinase related protein-2, VYDFFVWL), pR 60 and Concanavalin A (Con A). Spots representing single IFN- γ producing T cells were counted and the number of background spots was subtracted from all samples. There were many spots detected after the stimulation with Con A (data not shown) indicating a good condition of the used lymphocytes. For each experimental group of mice the number of spots/ 1×10^6 cells are illustrated in Figure 3, the standard deviation of ex vivo-stimulated triplicates are given. One hour after injection blood was taken from the tail vein and serum was prepared to determine the induction of systemic TNF- α and IL-6 using cytokine-specific ELISAs (Figure 4).

Example 3:

The combined injection of random sequences containing deoxy-Inosine and an TRP-2 peptide induces a strong immune response against the TRP-2 peptide which can be further enhanced by the co-application of poly-L-arginine (pR 60).

| | |
|---------------------------|---|
| Mice | C57Bl/6 (Harlan/Olac) |
| Peptide | TRP-2-peptide (VYDFFVWL), a MHC class I (H-2K ^b)-restricted epitope of mouse tyrosinase related protein-2 (Blom et al., 1997) was synthesized by standard solid phase F-moc synthesis, HPLC purified and analyzed by mass spectroscopy for purity. Dose: 300 μ g/mouse |
| Poly-L-arginine 60 (pR60) | Poly-L-arginine with an average degree of polymerization of 60 arginine residues; SIGMA chemicals Dose: 100 μ g/mouse |
| CpG-ODN 1668 | thiophosphate substituted ODNs containing a CpG motif: tcc atg <u>acg ttc</u> ctg atg ct, were synthesized by NAPS GmbH, Göttingen. |

00070

- 22 -

wdi thiophosphate substituted ODNs: nhh hhh wdi
nhh hhh hhh wn were synthesized by NAPS GmbH,
Göttingen.
Dose: 5nmol/mouse

wdidin thiophosphate substituted ODNs: nhh hhh wdi
nhh hhh hhh wn were synthesized by NAPS GmbH,
Göttingen.
Dose: 5nmol/mouse

wdid thiophosphate substituted ODNs: nhh hhh wdi
dhh hhh hhh wn were synthesized by NAPS GmbH,
Göttingen.
Dose: 5nmol/mouse

wdidid thiophosphate substituted ODNs: nhh wdi did
hhh hdi ddi dh were synthesized by NAPS GmbH,
Göttingen.
Dose: 5nmol/mouse

Experimental groups (5 mice per group)

1. TRP-2
2. TRP-2 + pR 60
3. TRP-2 + CpG 1668
4. TRP-2 + wdi
5. TRP-2 + wdidin
6. TRP-2 + wdid
7. TRP-2 + wdidid
8. TRP-2 + CpG 1668 + pR 60
9. TRP-2 + wdi + pR 60
10. TRP-2 + wdidin + pR 60
11. TRP-2 + wdid + pR 60
12. TRP-2 + wdidid + pR 60

On day 0 mice were injected into each hind footpad with a total volume of 100µl (50µl per footpad) containing the above mentioned compounds. Animals were sacrificed 4 days after injection and popliteal lymph nodes were harvested. Lymph nodes were passed through a 70µm cell strainer and washed twice with DMEM medium (GIBCO BRL) containing 5% fetal calf serum (FCS, SIGMA chemicals). Cells were adjusted to 2×10^6 cells/ml in DMEM/5% FCS.

033270

- 23 -

IFN- γ ELISPOT assay was carried out in triplicates as described (Miyahira et al., 1995). This method is a widely used procedure allowing the quantification of antigen-specific T cells. Lymphocytes were stimulated ex vivo in triplicates with medium (background), TRP-2-peptide, an irrelevant OVA₂₅₇₋₂₆₄-peptide, pR 60 and Concanavalin A (Con A). Spots representing single IFN- γ producing T cells were counted and the number of background spots was subtracted from all samples. There were many spots detected after the stimulation with Con A (data not shown) indicating a good condition of the used lymphocytes. For each experimental group of mice the number of spots/ 1×10^6 cells are illustrated in Figure 5, the standard deviation of ex vivo-stimulated triplicates are given.

03370

- 24 -

References

Andreu, D., and Rivas, L. (1998). Animal antimicrobial peptides: an overview. *Biopolymers* 47, 415-433.

Ballas, Z. K., Rasmussen, W. L., and Krieg, A. M. (1996). Induction of NK activity in murine and human cells by CpG motif in oligodeoxynucleotides and bacterial DNA. *J Immunol* 157, 1840-1845.

Bloom, B. R., and Widdus, R. (1998). Vaccine visions and their global impact. *Nat Med* 4, 480-484.

Bloom, M. B., Perry-Lalley, D., Robbins, P. F., Li, Y., el-Gamil, A., Rosenberg, S. A., and Yang, J. C. (1997). Identification of tyrosinase-related protein 2 as a tumor rejection antigen for the B 16 melanoma. *J Exp Med* 185, 453-459.

Buschle, M., Schmidt, W., Berger, M., Schaffner, G., Kurzbauer, R., Killisch, I., Tiedemarm, J.K., Trska, B., Kirlappos, H., Mechtler, K., Schilcher, F., Gabler, C., and Birnstiel, M. L. (1998). Chemically defined, cell-free cancer vaccines: use of tumor antigen-derived peptides or polypeptide proteins for vaccination. *Gene Ther. Mol. Biol.* 1, 309-321

Buschle, M., Schmidt, W., Zauner, W., Mechtler, K., Trska, B., Kirlappos, H., and Birnstiel, M.L. (1997). Transloading of tumor antigen-derived peptides into antigen-presenting cells. *Proc. Natl. Acad. Sci. USA* 94, 3256-3261

Cavanaugh, P.F., Jr., Ho, Y-K, and Bardos, T.J. (1996). The activation of murine macrophages and natural killer cells by the Partially thiolated double stranded RNA poly (1). mercapto poly(C). *Res.Comm.Mol.Pathol.Pharmacol.* 91, 131-147

Chace, J. H., Hooker, N. A., Mildenstein, K. L., Krieg, A. M., and Cowdery, J. S. (1997). Bacterial DNA-induced NK cell IFN-gamma production is dependent on macrophage secretion of IL- 12. *Clin Immunol Immunopathol* 84, 185-193.



- 25 -

Schorr, J., and Krieg, A. M. (1998). CpG DNA is a potent enhancer of specific immunity in mice immunized with recombinant hepatitis B surface antigen. *J Immunol* 160, 870-876.

Deng, G. M., Nilsson, I. M., Verdrengh, M., Collins, L. V., and Tarkowski, A. (1999). Intra-articularly localized bacterial DNA containing CpG motifs induces arthritis. *Nat Med* 5, 702-705.

Ganz, T. (1999). Defensins and host defense [comment]. *Science* 286, 420-421.

Ganz, T., and Lehrer, R. I. (1999). Antibiotic peptides from higher eukaryotes: biology and applications. *Mol Med Today* 5, 292-297.

Hancock, R. E. (1999). Host defence (cationic) peptides: what is their future clinical potential? *Drugs* 57, 469-473.

Harlow, E., and Lane, D. (1988). *Antibodies: a laboratory manual* (Cold Spring Harbor: Cold Spring Harbor Laboratory).

Hartmann, G., Weiner, G. J., and Krieg, A. M. (1999). CpG DNA: A potent signal for growth, activation, and maturation of human dendritic cells. *Proc Natl Acad Sci U S A* 96, 9305-9310.

Hoffmann, J. A., Kafatos, F. C., Janeway, C. A., and Ezekowitz, R. A. (1999). Phylogenetic perspectives in innate immunity. *Science* 284, 1313-1318.

Klinman, D. M., Yi, A. K., Beaucage, S. L., Conover, J., and Krieg, A. M. (1996). CpG motifs present in bacteria DNA rapidly induce lymphocytes to secrete interleukin 6, interleukin 12, and interferon gamma. *Proc Natl Acad Sci U S A* 93, 2879-2883.

Krieg, A. M. (1999). CpG DNA: a novel immunomodulator [letter]. *Trends Microbiol* 7, 64-5.

Krieg, A. M. (1996). An innate immune defense mechanism based on the recognition of CpG motifs in bacterial DNA. *Proc Natl Acad Sci U S A* 93, 2879-2883.

Krieg, A. M., Yi, A. K., Matson, S., Waldschmidt, T. J., Bishop, G. A., Teasdale, R., Koretzky, G. A., and Klinman, D. M. (1995). CpG motifs in bacterial DNA trigger direct B-cell activation. *Nature* 374, 546-549.

Krieg, A. M., Yi, A. K., Schorr, J., and Davis, H. L. (1998). The role of CpG dinucleotides in DNA vaccines. *Trends Microbiol* 6, 23-27.

Lethe, B., van den Eynde, B., van Pel, A., Corradin, G., and Boon, T. (1992). Mouse tumor rejection antigens P815A and P815B: two epitopes carried by a single peptide. *Eur J Immunol* 22, 2283-2288.

Liljeqvist, S., and Stahl, S. (1999). Production of recombinant subunit vaccines: protein immunogens, live delivery systems and nucleic acid vaccines. *J Biotechnol* 73, 1-33.

Lipford, G. B., Heeg, K., and Wagner, H. (1998). Bacterial DNA as immune cell activator. *Trends Microbiol* 6, 496-500.

Manetti, R., Annunziato, F., Tomasevic, L., Gianno, V., Parronchi, P., Romagnani, S. and Maggi, E. (1995). Polyinosinic acid: polycytidylic acid promotes T helper type 1-specific immune responses by stimulating macrophage production of interferon- α and interleukin-12. *Eur. J. Immunol.* 25, 2656-2660

Mosmann, T. R., Cherwinski, H., Bond, M. W., Giedlin, M. A., and Coffman, R. L. (1986). Two types of murine helper T cell clone. 1. Definition according to profiles of lymphokine activities and secreted proteins. *J Immunol* 136, 2348-2357.

Nossal, G. (1998). Living up to the legacy. *Nat Med* 4, 475-476

Oxenius, A., Martinic, M M., Hengartner, H., and Kleenerman, P. (1999). CpG-containing oligonucleotides are efficient adjuvants for induction of protective antiviral immune responses with T-cell peptide vaccines. *J Virol* 73, 4120-4126

03270

- 27 -

Paillard, F. (1999). CpG: the double-edged sword [comment]. *Hum Gene Ther* 10, 2089-2090.

Pamer, E. G., Harty, J. T., and Bevan, M. J. (1991). Precise prediction of a dominant class I MHC-restricted epitope of *Listeria monocytogenes*. *Nature* 353, 852-855.

Parronchi, P., Brugnolo, F., Annunziato, F., Manuelli, C., Sampognaro, S., Mavilia, C., Romagnani, S., and Maggi, E. (1999). Phosphorothioate oligodeoxynucleotides promote the in vitro development of human allergen-specific CD4+ T cells into Th1 effectors. *J Immunol* 163, 5946-5953.

Pisetsky, D. S. (1997). Immunostimulatory DNA: a clear and present danger? *Nat Med* 3, 829-831.

Pisetsky, D. S. (1999). The influence of base sequence on the immunostimulatory properties of DNA. *Immunol Res* 19, 35-46.


Rammensee, H.G., Friede, T., Stevanovic S. (1995), MHC ligands and peptide motifs: first listing. *Immunogenetics* 41, 178-228

Rodrigues, M., Nussenzweig, R. S., Romero, P., and Zavala, F. (1992). The in vivo cytotoxic activity of CD8+ T cell clones correlates with their levels of expression of adhesion molecules. *J Exp Med* 175, 895-905.

Roitt, I., Brostoff, J., and Male, D. (1998). *Immunology* (London: Mosby International Ltd).

Rotzschke, O., Falk, K., Stevanovic, S., Jung, G., Walden, P., and Rammensee, H. G. (1991). Exact prediction of a natural T cell epitope. *Eur J Immunol* 21, 2891-2894.

Schmidt, W., Buschle, M., Zauner, W., Kirlappos, H., Mechtler, K., Trska, B., and Bimstiel, M.L. (1997). Cell-free tumor antigen peptide-based cancer vaccines. *Proc. Natl. Acad. Sci. USA* 94, 3262-3267



K., and Krieg, A. M. (1997). CpG motifs in bacterial DNA cause inflammation in the lower respiratory tract, *J Clin Invest* 100, 68-73.

Shimonkevitz, R., Colon, S., Kappler, J. W., Marrack, P., and Grey, H. M. (1984). Antigen recognition by H2-restricted T cells 11. A tryptic ovalbumin peptide that substitutes for processed antigen. *J Immunol* 133, 2067-2074.

Simmaco, M., Mignogna, G., and Barra, D. (1998). Antimicrobial peptides from amphibian skin: what do they tell us? *Biopolymers* 47, 435-450.

Parbier, K., and Walden, P. (1999). T cell receptor specificity and mimotopes. *Curr Opin Immunol* 11, 214-218.

Sparwasser, T., Koch, E. S., Vabulas, R. M., Heeg, K., Lipford, G. B., Ellwart, J. W., and Wagner, H. (1998). Bacterial DNA and immunostimulatory CpG oligonucleotides trigger maturation and activation of murine dendritic cells. *Eur J Immunol* 28, 2045-2054.

Sparwasser, T., Miethke, T., Lipford, G., Borschert, K., Hacker, H., Heeg, K., and Wagner, H. (1997). Bacterial DNA causes septic shock [letter]. *Nature* 386, 336-337.

Sparwasser, T., Miethke, T., Lipford, G., Erdmann, A., Hacker, H., Heeg, K., and Wagner, H. (1997). Macrophages sense pathogens via DNA motif: induction of tumor necrosis factor- α -mediated shock. *Eur J Immunol* 27, 1671-1679.

Weiner, G. J., Liu, H. M., Wooldridge, J. E., Dahle, C. E., and Krieg, A. M. (1997). Immunostimulatory oligodeoxynucleotides containing the CpG motif are effective as immune adjuvants in tumor antigen immunization. *Proc Natl Acad Sci U S A* 94, 10833-10837.


Yew, N. S., Wang, K. X., Przybylska, M., Bagley, R. G., Stedman, M., Marshall, J., Scheule, R. K., and Cheng, S. H. (1999). Contribution of plasmid DNA to inflammation in the lung after ad-

030270

- 29 -

Claims:

1. Immunostimulatory oligodeoxynucleic acid molecule (ODN) having the structure according to formula (I)



(I),

any X is O or S,

wherein

any NMP is a 2' deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine-, deoxyguanosine-, deoxyinosine-, deoxycytosine-, deoxyuridine-, deoxythymidine-, 2-methyl-deoxyinosine-, 5-methyl-deoxycytosine-, deoxypseudouridine-, deoxyribosepurine-, 2-amino-deoxyribosepurine-, 6-S-deoxyguanine-, 2-dimethyl-deoxyguanosine- or N-isopentenyl-deoxyadenosine-monophosphate or -monothiophosphat,

NUC is a 2' deoxynucleoside, selected from the group consisting of deoxyadenosine-, deoxyguanosine-, deoxyinosine-, deoxycytosine-, deoxyuridine-, deoxythymidine-, 2-methyl-deoxyinosine-, 5-methyl-deoxycytosine-, deoxypseudouridine-, deoxyribosepurine-, 2-amino-deoxyribosepurine-, 6-S-deoxyguanine-, 2-dimethyl-deoxyguanosine- or N-isopentenyl-deoxyadenosine,

a and b are integers from 0 to 100 with the proviso that a + b is between 4 and 150,

B and E are common groups for 5' or 3' ends of nucleic acid molecules.

2. ODN according to claim 1, wherein any NMP is selected from

03370

- 30 -

nosine-, deoxycytosine-, deoxyuridine-, deoxythymidine-, 2-methyl-deoxyinosine-, 5-methyl-deoxycytosine-monophosphate or -monothiophosphate.

3. ODN according to claim 1 or 2, characterized in that $a + b$ is between 10 and 60, preferably between 15 and 40.

4. ODN according to any of claims 1 to 3, characterized in that at least one of X_1 and X_2 is S and at least one of X_3 and X_4 is O and preferably any NMP is a nucleoside-monothiophosphate.

5. ODN according to any of claims 1 to 4, characterized in that it contains the sequence

nhh hhh wdi nhh hhh hhh wn,
nhh wdi din hhh hdi ndi nh,
nhh hhh wdi dhh hhh hhh wn or
nhh wdi did hhh hdi ddi dh,

wherein

any n is a 2'-deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine-, deoxyguanosine-, deoxycytosine- or deoxythymidine-monophosphate or -monothiophosphate,

any h is a 2'-deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine-, deoxycytosine- or deoxythymidine-monophosphate or -monothiophosphate

i is deoxyinosine-monophosphate or -monothiophosphate,

any w is a 2'-deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine- or deoxythymidine-monophosphate or -monothiophosphate, and

any d is a 2'-deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine-, deoxyguanosine- or deoxythymidine-monophosphate or -monothiophosphate.

6. ODN according to any one of claims 1 to 5, characterized in that it contains at least one 2'-deoxycytosine-monophosphate or -monothiophosphate 3'-adjacent to a 2'-deoxyinosine-monophosphate or -monothiophosphate.

7. ODN according to any one of claims 1 to 6, characterized in

033770

- 31 -

gacitt,
iacitt,
gaicct,
iaicct,

wherein

a is deoxyadenosine-monophosphate or -monothiophosphate,
g is deoxyguanosine-monophosphate or -monothiophosphate,
i is deoxyinosine-monophosphate or -monothiophosphate,
c is deoxycytosine-monophosphate or -monothiophosphate and
t is deoxythymidine-monophosphate or -monothiophosphate.

8. ODN according to any one of claims 1 to 7, characterized in that it contains the sequence

w di,
w did,
w didin or,
w didid,

wherein w, d, i and n are defined as above.

9. ODN according to any one of claims 1 to 8, characterized in that B and E are selected independently from the group consisting of -H, -CH₃, -COH, -COCH₃, -OH, -CHO, -PO₄, -PSO₃, -PS₂O₂, -PS₃O, -PS₄, -SO₃, -PO₄-(CH₂)₁₋₆-NH₂ or -PO₄-(CH₂)₁₋₆-NH-Label.

10. Use of an ODN according to any one of claims 1 to 9 as a medicine, especially as an immunostimulatory agent.

11. Pharmaceutical composition comprising an ODN according to any one of claims 1 to 9.

12. Pharmaceutical composition comprising

- an ODN according to any one of claims 1 to 9 and
- an antigen.

13. Pharmaceutical composition according to claim 11 or 12, characterized in that it further comprises a polycationic polymer, preferably a polycationic peptide, especially polyarginine, polylysine or an antimicrobial peptide, especially a cathelicidin derived antimicrobial peptide or a growth hormone espe-

03070

- 32 -

14. Pharmaceutical composition according to any one of claims 11 to 13 further comprising further active ingredients, especially cytokines, antiinflammatory substances, antimicrobial substances or combinations thereof.

15. Pharmaceutical composition according to any one of claims 11 to 14, characterized in that it further contains auxiliary substances, especially a pharmaceutically acceptable carrier, buffer substances, stabilizers or combinations thereof.

16. Pharmaceutical composition according to any one of claims 11 to 15, characterized in that it contains 1 ng to 1 g, preferably 100 ng to 10 mg, especially 10 mg to 1 mg, of one or more ODNs according to any one of claims 1 to 9.

17. Use of an ODN according to any one of claims 1 to 9 for the preparation of a vaccine.


DA/R

033270

- 33 -

ABSTRACT

Described is an immunostimulatory oligodeoxynucleic acid molecule (ODN) having the structure according to formula (I)



(I),

wherein

any NMP is a 2' deoxynucleoside monophosphate or monothiophosphate, selected from the group consisting of deoxyadenosine-, deoxyguanosine-, deoxyinosine-, deoxycytosine-, deoxyuridine-, deoxythymidine-, 2-methyl-deoxyinosine-, 5-methyl-deoxycytosine-, deoxypseudouridine-, deoxyribosepurine-, 2-amino-deoxyribosepurine-, 6-S-deoxyguanine-, 2-dimethyl-deoxyguanosine- or N-isopentenyl-deoxyadenosine-monophosphate or -monothiophosphate, NUC is a 2' deoxynucleoside, selected from the group consisting of deoxyadenosine-, deoxyguanosine-, deoxyinosine-, deoxycytosine-, deoxyuridine-, deoxythymidine-, 2-methyl-deoxyinosine-, 5-methyl-deoxycytosine-, deoxypseudouridine-, deoxyribosepurine-, 2-amino-deoxyribosepurine-, 6-S-deoxyguanine-, 2-dimethyl-deoxyguanosine- or N-isopentenyl-deoxyadenosine, any X is O or S,

a and b are integers from 0 to 100 with the proviso that a + b is between 4 and 150,

B and E are common groups for 5' or 3' ends of nucleic acid molecules,

as well as a pharmaceutical composition containing such ODNs.

A1973/2000

1/5

urtext

IFN- γ ELISPOT
day 4 after vaccination




Fig. 1
injection of OVA₂₅₇₋₂₆₄




Fig 2

Fig 3





Fig 4



A1973/2000

5/5

Untext

Fig 5

